



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
 General Certificate of Education  
 Advanced Subsidiary Level and Advanced Level

CANDIDATE  
 NAME

CENTRE  
 NUMBER

--	--	--	--

CANDIDATE  
 NUMBER

--	--	--	--



**CHEMISTRY**

**9701/34**

Paper 32 Advanced Practical Skills

**May/June 2010**

**2 hours**

Candidates answer on the Question Paper.

Additional Materials: As listed in the Instructions to Supervisors

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.  
 Give details of the practical session and laboratory where appropriate, in the boxes provided.  
 Write in dark blue or black pen.  
 You may use a soft pencil for any diagrams, graphs or rough working.  
 Do not use staples, paper clips, highlighters, glue or correction fluid.  
 DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.  
 You may lose marks if you do not show your working or if you do not use appropriate units.  
 Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.

At the end of the examination, fasten all your work securely together.  
 The number of marks is given in brackets [ ] at the end of each question or part question.

<b>Session</b>
<b>Laboratory</b>

For Examiner's Use	
<b>1</b>	
<b>2</b>	
<b>Total</b>	

This document consists of **11** printed pages and **1** blank page.

**1 Read through question 1 before starting any practical work.**For  
Examiner's  
Use

You are provided with the following reagents.

- weighing bottles/tubes labelled **FB 1**, **FB 2** and **FB 3**; each containing a different mass of sodium hydrogencarbonate,  $\text{NaHCO}_3$
- additional solid sodium hydrogencarbonate (approximately 10g)
- **FB 4**,  $3.0 \text{ mol dm}^{-3}$  hydrochloric acid,  $\text{HCl}$

The reaction of sodium hydrogencarbonate with hydrochloric acid is endothermic.

By measuring the maximum temperature decrease when the different masses of sodium hydrogencarbonate react with hydrochloric acid you are to determine the enthalpy change of neutralisation for 1 mol of  $\text{NaHCO}_3$  with  $\text{HCl}$ .

**(a) Method**

- Weigh the bottle/tube containing the sodium hydrogencarbonate labelled **FB 1**.
- Support the plastic cup in the  $250 \text{ cm}^3$  beaker.
- Use the measuring cylinder to transfer  $30 \text{ cm}^3$  of **FB 4** into the plastic cup.
- Place the thermometer in the acid in the plastic cup and record the steady temperature of the acid.
- Add the contents of the weighed tube, **FB 1**, to the acid in the plastic cup, a little at a time with constant stirring.
- You should add the solid as quickly as possible – taking care to minimise any acid spray from the plastic cup.  
*Avoid breathing any fumes from the experiment.*
- Record the minimum temperature obtained in the reaction.
- Reweigh the emptied tube, **FB 1**, containing any remaining solid that was not tipped from the tube.
- Empty and rinse the plastic cup. Rinse the thermometer. Shake dry the plastic cup.
- Repeat the experiment using tubes labelled **FB 2** and **FB 3**. In each experiment use  $30 \text{ cm}^3$  of **FB 4**.

Carry out **two further experiments**.

Using the empty weighing bottles/tubes, labelled **FB 5** and **FB 6**, weigh two further masses of sodium hydrogencarbonate. Choose masses to enable you to plot an appropriate graph of temperature change against mass of sodium hydrogencarbonate.

**Results**


Record your results in an appropriate form showing, for each experiment, the measurements of mass and temperature and the calculated temperature fall.

i	
ii	
iii	
iv	
v	
vi	
vii	
viii	
ix	

[9]

- (b) Use the grid below to plot a graph of decrease in temperature (*y-axis*) against the mass of sodium hydrogencarbonate added (*x-axis*).  
Draw a line of best fit through the plotted points.  
You should consider if the best-fit line passes through the origin (0,0) of the graph.

For  
Examiner's  
Use



i	
ii	
iii	
iv	

[4]

- (c) Explain why the mass of  $\text{NaHCO}_3$  is plotted on the *x-axis* rather than on the *y-axis*.

.....  
 ..... [1]

- (d) Construct the balanced equation for the reaction of  $\text{NaHCO}_3$  with hydrochloric acid.

..... [1]

- (e) Calculate the gradient of your graph.

Show all of your working clearly, both construction lines on the graph and working in the calculation.

[3]

- (f) Although there is insufficient acid in  $30\text{ cm}^3$  of **FB 4** to neutralise 1 mol of  $\text{NaHCO}_3$  it is possible to calculate the theoretical fall in temperature for this reaction.

Use your answer from (e) to calculate this value.

[ $A_r$ : C, 12.0; H, 1.0; Na, 23.0; O, 16.0]

The theoretical fall in temperature for 1 mol of  $\text{NaHCO}_3$  = ..... °C [1]

- (g) Calculate the theoretical enthalpy change for the neutralisation of 1 mol of  $\text{NaHCO}_3$  by hydrochloric acid. Give your answer in  $\text{kJ mol}^{-1}$  and include the correct sign for the reaction.

[4.3J are absorbed or released when the temperature of  $1\text{ cm}^3$  of solution changes by  $1\text{ }^\circ\text{C}$ .]

$\Delta H =$  .....  $\text{kJ mol}^{-1}$  [2]

- (h) Suggest **two** ways in which your apparatus could be modified to reduce transfer of heat from the surroundings to the solution in the apparatus.

modification 1 .....

.....

modification 2 .....

..... [1]

- (i) State and explain why the experiment would be more accurate if the volumes of **FB 4** had been measured using a burette instead of a measuring cylinder.

.....

..... [1]

- (j) The mass of  $\text{NaHCO}_3$  used in a further experiment and its associated temperature change are shown in the tables below.

- The mass was obtained on a balance reading to 1 decimal place.
- The thermometer used was graduated at 1 °C.

Complete the table to show the errors in these results.

mass of $\text{NaHCO}_3$	5.6 g	temperature change	-12.0 °C
maximum error in a single balance reading	± ..... g	maximum error in a single thermometer reading	± ..... °C
% error in measured mass	..... %	% error in temperature change	..... %

[2]

- (k) Two students add 6.0 g of sodium carbonate to 50.00 cm<sup>3</sup> of 2.0 mol dm<sup>-3</sup> hydrochloric acid. Each student repeats the experiment a number of times. The thermometer readings and temperature changes obtained consistently by each student are shown below.

	initial temperature / °C	final temperature / °C	temperature rise / °C
student 1	20.0	28.0	8.0
student 2	19.0	27.0	8.0

Suggest the type of error shown by these results.

..... [1]

[Total: 26]

- 2 **FB 7** and **FB 8** are aqueous solutions of salts. One of these contains **two** cations and one anion.  
The other contains one cation and one anion.  
Both **FB 7** and **FB 8** have a common cation.

You will carry out tests to deduce the following.

- the cations present in each solution
- whether a sulfate ion is present in either solution

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate and the colour of the precipitate

Where gases are released they should be identified by a test, **described in the appropriate place in your observations.**

You should indicate clearly at what stage in a test a change occurs.

Marks are **not** given for chemical equations.

**No additional tests for ions present should be attempted.**

**If any solution is warmed directly with a Bunsen burner a boiling-tube MUST be used.**

Rinse and reuse test-tubes where possible.

- (a) Use information from the Qualitative Analysis Notes on page 11 to select a pair of reagents that, **used together**, determine whether a sulfate ion is present in either solution.

The reagents are .....

followed by .....

[1]

- (b) Use your chosen reagents to carry out tests on **FB 7** and **FB 8**.  
Record your results in an appropriate form in the space below.

[2]

- (c) From your observations in (b) show with a tick which of the following statements is true.

For  
Examiner's  
Use

FB 7 contains the sulfate ion	
FB 8 contains the sulfate ion	
neither solution contains the sulfate ion	

Explain the evidence that supports your conclusion.

.....  
..... [1]

- (d) Carry out the following tests on the solutions **FB 7** and **FB 8**.  
Record your observations below.

<i>test</i>	<i>observations</i>	
	<b>FB 7</b>	<b>FB 8</b>
To 1 cm depth of solution in a boiling-tube, add 2 cm depth of aqueous sodium hydroxide; then		
warm the solution gently.  <b>Care is needed when heating aqueous sodium hydroxide.</b>		
To 1 cm depth of solution in a test-tube, add 2 cm depth of aqueous ammonia.		

[3]

- (e) To 1 cm depth of **FB 7** in a test-tube add 1 cm depth of sodium hydroxide.  
Leave to stand for a few minutes.

observation

.....  
..... [1]

- (f) From your observations in (d) and (e) you should be able to identify the common cation in the solutions and the second cation in **one** of the solutions.

The common cation present in both solutions is .....

The second cation contained in one of the solutions is .....

Explain how your observations support your conclusions for

- (i) the common cation,

.....  
.....

- (ii) the second cation.

.....  
.....

[1]

**Read through the remainder of question 2 before starting further practical work.**

**Heat a half-full 250 cm<sup>3</sup> beaker of water for use as a hot water-bath.**

- (g) **FB 9, FB 10, FB 11** and **FB 12** are organic compounds. Each contains one of the following different functional groups.

- primary alcohol
- secondary alcohol
- aldehyde
- ketone

You are to react each of these compounds with some of the following reagents.

- acidified aqueous potassium dichromate(VI)
- 2,4-dinitrophenylhydrazine (2,4-DNPH) reagent
- ammoniacal silver nitrate (Tollens' reagent)

You are provided with the first two reagents. You must prepare the last of these reagents, Tollens' reagent, immediately before use. Follow the instructions in the box below.

To 2 cm depth of aqueous silver nitrate in a boiling-tube add ½ cm depth of aqueous sodium hydroxide. This will produce a brown precipitate of silver(I) oxide. Add aqueous ammonia a little at a time, with continuous shaking, until the brown precipitate **just** dissolves. **Do not add an excess of aqueous ammonia.**



In each of the following tests add a few drops of the reagent to 1 cm depth of **FB 9**, **FB 10**, **FB 11** and **FB 12** in separate test-tubes.

In the tests using acidified potassium dichromate(VI) and Tollens' reagent, if no initial reaction is seen, warm that tube and its contents in your hot water-bath. There is no need to heat any tube to which you have added 2,4-DNPH reagent.

Do **not** heat any tube with a naked flame.

Record your results in the table below.

Do **not** carry out tests for the shaded boxes.

For  
Examiner's  
Use

reagent	observations			
	FB 9	FB 10	FB 11	FB 12
acidified potassium dichromate(VI)				
2,4-DNPH reagent				
Tollens' reagent				

[3]

- (h) State which of the solutions contain alcohols. Explain the observations leading to your conclusion.

**FB** ..... and **FB** ..... contain alcohols.

explanation .....

.....

State which solution contains the ketone. Explain the observations leading to your conclusion.

**FB** ..... contains the ketone.

explanation .....

.....

[2]

[Total: 14]

Key: [ ppt. = precipitate. ]

## 1 Reactions of aqueous cations

ion	reaction with	
	NaOH(aq)	NH <sub>3</sub> (aq)
aluminium, Al <sup>3+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH <sub>4</sub> <sup>+</sup> (aq)	no ppt. ammonia produced on heating	–
barium, Ba <sup>2+</sup> (aq)	no ppt. (if reagents are pure)	no ppt.
calcium, Ca <sup>2+</sup> (aq)	white ppt. with high [Ca <sup>2+</sup> (aq)]	no ppt.
chromium(III), Cr <sup>3+</sup> (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu <sup>2+</sup> (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe <sup>2+</sup> (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe <sup>3+</sup> (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
lead(II), Pb <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
magnesium, Mg <sup>2+</sup> (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn <sup>2+</sup> (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. soluble in excess

[Lead(II) ions can be distinguished from aluminium ions by the insolubility of lead(II) chloride.]

## 2 Reactions of anions

<i>ion</i>	<i>reaction</i>
carbonate, $\text{CO}_3^{2-}$	$\text{CO}_2$ liberated by dilute acids
chromate(VI), $\text{CrO}_4^{2-}$ (aq)	yellow solution turns orange with $\text{H}^+$ (aq); gives yellow ppt. with $\text{Ba}^{2+}$ (aq); gives bright yellow ppt. with $\text{Pb}^{2+}$ (aq)
chloride, $\text{Cl}^-$ (aq)	gives white ppt. with $\text{Ag}^+$ (aq) (soluble in $\text{NH}_3$ (aq)); gives white ppt. with $\text{Pb}^{2+}$ (aq)
bromide, $\text{Br}^-$ (aq)	gives cream ppt. with $\text{Ag}^+$ (aq) (partially soluble in $\text{NH}_3$ (aq)); gives white ppt. with $\text{Pb}^{2+}$ (aq)
iodide, $\text{I}^-$ (aq)	gives yellow ppt. with $\text{Ag}^+$ (aq) (insoluble in $\text{NH}_3$ (aq)); gives yellow ppt. with $\text{Pb}^{2+}$ (aq)
nitrate, $\text{NO}_3^-$ (aq)	$\text{NH}_3$ liberated on heating with $\text{OH}^-$ (aq) and Al foil
nitrite, $\text{NO}_2^-$ (aq)	$\text{NH}_3$ liberated on heating with $\text{OH}^-$ (aq) and Al foil; NO liberated by dilute acids (colourless NO $\rightarrow$ (pale) brown $\text{NO}_2$ in air)
sulfate, $\text{SO}_4^{2-}$ (aq)	gives white ppt. with $\text{Ba}^{2+}$ (aq) or with $\text{Pb}^{2+}$ (aq) (insoluble in excess dilute strong acids);
sulfite, $\text{SO}_3^{2-}$ (aq)	$\text{SO}_2$ liberated with dilute acids; gives white ppt. with $\text{Ba}^{2+}$ (aq) (soluble in excess dilute strong acids)

## 3 Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia, $\text{NH}_3$	turns damp red litmus paper blue
carbon dioxide, $\text{CO}_2$	gives a white ppt. with limewater (ppt. dissolves with excess $\text{CO}_2$ )
chlorine, $\text{Cl}_2$	bleaches damp litmus paper
hydrogen, $\text{H}_2$	"pops" with a lighted splint
oxygen, $\text{O}_2$	relights a glowing splint
sulfur dioxide, $\text{SO}_2$	turns acidified aqueous potassium dichromate(VI) from orange to green

**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.