



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
 General Certificate of Education  
 Advanced Subsidiary Level and Advanced Level

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 1 4 1 7 9 8 4 1 1 0 \*

**CHEMISTRY**

**9701/35**

Advanced Practical Skills 1

**October/November 2013**

**2 hours**

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.  
 Give details of the practical session and laboratory where appropriate, in the boxes provided.  
 Write in dark blue or black pen.  
 You may use a soft pencil for any diagrams, graphs or rough working.  
 Do not use staples, paper clips, highlighters, glue or correction fluid.  
**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.  
 Electronic calculators may be used.  
 You may lose marks if you do not show your working or if you do not use appropriate units.  
 Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.

At the end of the examination, fasten all your work securely together.  
 The number of marks is given in brackets [ ] at the end of each question or part question.

<b>Session</b>
<b>Laboratory</b>

<b>For Examiner's Use</b>	
<b>1</b>	
<b>2</b>	
<b>3</b>	
<b>Total</b>	

This document consists of **11** printed pages and **1** blank page.

1 You are provided with a solution of an organic acid which is known to be one of the following.

- methanoic acid, HCOOH  
 propanoic acid, C<sub>2</sub>H<sub>5</sub>COOH  
 pent-2-enoic acid, CH<sub>3</sub>CH<sub>2</sub>CH=CHCOOH

The solution was made by dissolving 1.85 g of acid in 250 cm<sup>3</sup> of solution.

You are to suggest the identity of the acid by finding its relative molecular mass,  $M_r$ , using a titration method.

**FA 1** is the solution of the unknown organic acid.

**FA 2** is 0.100 mol dm<sup>-3</sup> sodium hydroxide, NaOH.

phenolphthalein indicator

**(a) Method**

- Fill the burette with **FA 2**.
- Pipette 25.0 cm<sup>3</sup> of **FA 1** into a conical flask.
- Titrate **FA 1** with **FA 2** using phenolphthalein as indicator.
- Perform a rough titration and record your burette readings in the space below.

The rough titre is ..... cm<sup>3</sup>.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record, in a suitable form below, all of your burette readings and the volume of **FA 2** added in each accurate titration.

For  
Examiner's  
Use

I	
II	
III	
IV	
V	
VI	
VII	

[7]

- (b) From your accurate titration results, obtain a suitable value to be used in your calculations. Show clearly how you obtained this value.

25.0 cm<sup>3</sup> of **FA 1** required ..... cm<sup>3</sup> of **FA 2** [1]

(c) **Calculations**

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- (i) Calculate the number of moles of sodium hydroxide in the volume of **FA 2** you have calculated in (b).

moles of NaOH = ..... mol

- (ii) One mole of any of the organic acids reacts with one mole of sodium hydroxide. Calculate the concentration, in mol dm<sup>-3</sup>, of the acid in **FA 1**.

concentration of the acid in **FA 1** = ..... mol dm<sup>-3</sup>

- (iii) Calculate the concentration, in g dm<sup>-3</sup>, of the acid used to make solution **FA 1**.

concentration of the acid in **FA 1** = ..... g dm<sup>-3</sup>

- (iv) Using your answers to (ii) and (iii), calculate the relative molecular mass,  $M_r$ , of the acid in **FA 1**.

$M_r$  of the acid = .....

- (v) Suggest which of the acids, methanoic, propanoic or pent-2-enoic acid, is present in FA 1.

Acid present is .....

- (vi) Suggest a test that could be carried out to distinguish pent-2-enoic acid from methanoic acid and propanoic acid.  
Give the expected result of your test.

test .....

expected result .....

[5]

[Total: 13]

- 2 The formula of hydrated iron(II) sulfate is  $\text{FeSO}_4 \cdot x\text{H}_2\text{O}$  where  $x$  shows the number of moles of water of crystallisation.

The value of  $x$  can be found by heating solid hydrated iron(II) sulfate to remove the water of crystallisation.

**FA 3** is hydrated iron(II) sulfate,  $\text{FeSO}_4 \cdot x\text{H}_2\text{O}$ .

**(a) Method**

Record **all** weighings, in an appropriate form, in the space below.

- Weigh and record the mass of the empty crucible.
- Tip the contents of the tube labelled **FA 3** into the weighed crucible. Reweigh and record the mass of the crucible and **FA 3**.
- Use a pipe-clay triangle to support the crucible and contents on a tripod.
- Heat **gently** for about three minutes.
- Leave the crucible to cool for approximately five minutes.

**While you are waiting for the crucible to cool, start work on Question 3.**

- When cool, reweigh the crucible with the residue.
- Reheat **gently** for three minutes, cool and reweigh the crucible until you are satisfied that all the water of crystallisation has been lost. It should not be necessary to reheat the crucible more than three times.

For  
Examiner's  
Use

I	
II	
III	
IV	
V	
VI	

[6]

- (b) (i) Calculate the mass of water lost and the mass of anhydrous iron(II) sulfate that remains after the heating process.

mass of water lost = ..... g

mass of anhydrous iron(II) sulfate = ..... g

- (ii) Determine the value of  $x$  in the formula of hydrated iron(II) sulfate,  $\text{FeSO}_4 \cdot x\text{H}_2\text{O}$ .  
( $A_r$ : H, 1.0; O, 16.0; S, 32.1; Fe, 55.8)

value of  $x$  = .....

[3]

- (c) A group of students carried out this practical and made their measurements correctly. The students calculated a value of 9 for  $x$ . The textbook value of  $x$  is less than 9.

- (i) Suggest an error in the practical procedure of the experiment that could account for this result and explain why this gives a value of  $x$  that is too high.

.....  
 .....  
 .....  
 .....

- (ii) Suggest a modification that could be made to the experimental procedure to reduce this error. Explain why this modification should give an answer for  $x$  that is closer to the textbook value.

.....  
 .....  
 .....

[3]

[Total: 12]

### 3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations.**

You should indicate clearly at what stage in a test a change occurs.

Marks are **not** given for chemical equations.

**No additional tests for ions present should be attempted.**

**If any solution is warmed, a boiling tube MUST be used.**

Rinse and reuse test-tubes and boiling tubes where possible.

**Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.**

(a) You are provided with a solid **FA 4**. **FA 4** is a mixture that contains two cations and two anions.

(i) Place a spatula measure of **FA 4** in a **hard-glass** test-tube. Heat the solid and identify the gas given off. Record all your observations.

.....

.....

.....

(ii) To a spatula measure of **FA 4** in a test-tube, add a 1 cm depth of dilute nitric acid. Record your observations.

.....

.....

- (iii) To a spatula measure of **FA 4** in a test-tube, add approximately a 2 cm depth of distilled water to make a solution. Divide the solution into two portions.

To the first portion, add a 1 cm depth of aqueous sodium hydroxide. Record your observations.

.....  
 .....

To the second portion, add a few drops of aqueous silver nitrate, then add a 1 cm depth of dilute nitric acid. Record your observations.

.....  
 .....

- (iv) Use your results from (i) to (iii) to identify two anions and one cation that are present in **FA 4**.

anions present ..... and .....

cation present .....

- (v) What further test could be carried out on **FA 4** to confirm the presence of the cation you suggested in (iv)? You should state the reagent to be used and the expected result.

**Do not carry out this test.**

.....  
 .....

- (vi) To a spatula measure of **FA 4** in a test-tube, add a 1 cm depth of distilled water to make a solution. To this solution, add a few drops of aqueous barium chloride or barium nitrate.

Describe the appearance of the precipitate formed and state its identity.

appearance of precipitate .....

identity of precipitate .....

[9]



(b) **FA 5**, **FA 6**, **FA 7** and **FA 8** are aqueous solutions each containing one of the ions  $Al^{3+}$ ,  $Ca^{2+}$ ,  $Zn^{2+}$  and  $Pb^{2+}$ .

For  
Examiner's  
Use

(i) Carry out the following tests. Record your observations in the spaces provided in the table.

<i>test</i>	<i>observations</i>			
	<b>FA 5</b>	<b>FA 6</b>	<b>FA 7</b>	<b>FA 8</b>
To a 1 cm depth of solution in a test-tube, add a few drops of aqueous sodium hydroxide, then				
add excess aqueous sodium hydroxide.				
To a 1 cm depth of solution in a test-tube, add a few drops of aqueous ammonia, then				
add excess aqueous ammonia.				
To a 1 cm depth of solution in a test-tube, add a 1 cm depth of aqueous potassium iodide.				

(ii) Use the Qualitative Analysis Notes on page **10** to identify the cation present in each of the solutions.

**FA 5** is ..... , **FA 6** is ..... , **FA 7** is ..... , **FA 8** is .....

[6]

[Total: 15]

## Qualitative Analysis Notes

Key: [ppt. = precipitate]

## 1 Reactions of aqueous cations

ion	reaction with	
	NaOH(aq)	NH <sub>3</sub> (aq)
aluminium, Al <sup>3+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH <sub>4</sub> <sup>+</sup> (aq)	no ppt. ammonia produced on heating	–
barium, Ba <sup>2+</sup> (aq)	no ppt. (if reagents are pure)	no ppt.
calcium, Ca <sup>2+</sup> (aq)	white ppt. with high [Ca <sup>2+</sup> (aq)]	no ppt.
chromium(III), Cr <sup>3+</sup> (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu <sup>2+</sup> (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe <sup>2+</sup> (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe <sup>3+</sup> (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
lead(II), Pb <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
magnesium, Mg <sup>2+</sup> (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn <sup>2+</sup> (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. soluble in excess

[Lead(II) ions can be distinguished from aluminium ions by the insolubility of lead(II) chloride.]

## 2 Reactions of anions

<i>ion</i>	<i>reaction</i>
carbonate, $\text{CO}_3^{2-}$	$\text{CO}_2$ liberated by dilute acids
chromate(VI), $\text{CrO}_4^{2-}(\text{aq})$	yellow solution turns orange with $\text{H}^+(\text{aq})$ ; gives yellow ppt. with $\text{Ba}^{2+}(\text{aq})$ ; gives bright yellow ppt. with $\text{Pb}^{2+}(\text{aq})$
chloride, $\text{Cl}^-(\text{aq})$	gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$ ); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$
bromide, $\text{Br}^-(\text{aq})$	gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$ ); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$
iodide, $\text{I}^-(\text{aq})$	gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$ ); gives yellow ppt. with $\text{Pb}^{2+}(\text{aq})$
nitrate, $\text{NO}_3^-(\text{aq})$	$\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and Al foil
nitrite, $\text{NO}_2^-(\text{aq})$	$\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and Al foil; $\text{NO}$ liberated by dilute acids (colourless $\text{NO} \rightarrow$ (pale) brown $\text{NO}_2$ in air)
sulfate, $\text{SO}_4^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ or with $\text{Pb}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids)
sulfite, $\text{SO}_3^{2-}(\text{aq})$	$\text{SO}_2$ liberated with dilute acids; gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acids)

## 3 Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia, $\text{NH}_3$	turns damp red litmus paper blue
carbon dioxide, $\text{CO}_2$	gives a white ppt. with limewater (ppt. dissolves with excess $\text{CO}_2$ )
chlorine, $\text{Cl}_2$	bleaches damp litmus paper
hydrogen, $\text{H}_2$	“pops” with a lighted splint
oxygen, $\text{O}_2$	relights a glowing splint
sulfur dioxide, $\text{SO}_2$	turns acidified aqueous potassium dichromate(VI) from orange to green

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.