



GCE MARKING SCHEME

**CHEMISTRY
AS/Advanced**

SUMMER 2012

CH1
Section A

1.  [1]

2. $1/12^{\text{th}}$ mass of one atom of carbon-12. [1]

3. C [1]

4. (a)

C	O	Cl	
<u>12.1</u>	<u>16.2</u>	<u>71.7</u>	(1)
12	16	35.5	
1.01	1.01	2.02	
1	1	2	

Formula = COCl_2 (1) [2]

(b) M_r / molecular mass / number of atoms of any element in compound [1]

5. (a) **C B D E A** [2]
(1 mark if one mistake e.g. **A** in wrong place)

(b) **Z** (1)
Si is in Group 4 therefore large jump in ionisation energy would be after the fourth ionisation, not before it / **W**, **X** and **Y** have a large jump before the fourth ionisation energy so cannot be in Group 4 (1)

[2]

Total [10]

Section B

6. (a) (i) 12 [1]
- (ii) 14 [1]
- (iii) Percentage / abundance / ratio / proportion of each isotope [1]
- (b) (i) 0.125 g [1]
- (ii) e.g. Cobalt-60 (1) in radiotherapy (1) / Carbon-14 (1) in radio carbon dating (1) / Iodine-131 (1) as a tracer in thyroid glands (1) [2]
- (c) (i) Atoms are hit by an electron beam / electrons fired from an electron gun (and lose electrons) [1]
- (ii) To be able to accelerate the ions (to high speed) / so that they can be deflected by a magnetic field
- no credit for 'so that *atoms* can be deflected...'
[1]
- (iii) They are deflected by a magnetic field / according to the m/z ratio [1]
- (d)
- | | | | | |
|----|----|----------|----|----|
| 1s | 2s | 2p | 3s | 3p |
| ↓↑ | ↓↑ | ↓↑ ↓↑ ↓↑ | | |
- [1]
- (e) (i) $\text{Mg}_3\text{N}_2 + 6\text{H}_2\text{O} \longrightarrow 3\text{Mg}(\text{OH})_2 + 2\text{NH}_3$ [1]
- (ii) moles $\text{Mg}(\text{OH})_2 = 1.75/58.32 = 0.0300$ (1)
- moles $\text{Mg}_3\text{N}_2 = 0.0100$ (1)
- mass $\text{Mg}_3\text{N}_2 = 0.01 \times 100.9 = 1.01$ g (1) [3]
- must be **3 significant figures** to gain third mark

Total [14]


7. (a) Plotting (2)
 Best fit line (1) [3]
- (b) (i) C (1)
 Curve steeper (1) [2]
- (ii) Concentration of acid is greatest [1]
- (c) $44 \text{ cm}^3 (\pm 1 \text{ cm}^3)$ [1]
- (d) Moles Mg = $0.101/24.3 = 0.00416$ (1)
 Moles HCl = $2 \times 0.02 = 0.04$ (1) [2]
- (e) (i) Mg is not the limiting factor /
 Mg now in excess / HCl not in excess [1]
- (ii) Moles acid = $0.5 \times 0.04 = 0.02$ (1)
 Volume $\text{H}_2 = 0.01 \times 24 = 0.24 \text{ dm}^3$
 - correct unit needed (1) [2]
- (f) Lower the temperature of the acid (1)
 Reactants collide with less energy (1)
 Fewer molecules that have the required activation energy (1) [3]
 or Use pieces of magnesium (1) less surface area (1) less chance
 of successful collisions (1)
- QWC Selection of a form and style of writing appropriate to purpose
 and to complexity of subject matter. [1]

Total [16]

8. (a) Oil is non-renewable / will run out (1)
 Contribution of CO₂ to global warming (1)
 Oil has other important uses (1) [2]
 (Maximum 2 marks)
- (b) (i) Power stations / fossil fuels used to generate the electricity needed to make H₂ (1)
 Resulting in CO₂ formation (global warming) / acid rain (1)
 Manufacture of car produces pollution (1) [2]
 (Maximum 2 marks)
- QWC Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning [1]
- (ii) Disagree, no fuel is 100% safe / petrol can burn explosively
 (Accept agree if valid reason given e.g. in terms of lives being lost) [1]
- (c) (i) Hydrogen since frequency is inversely proportional to wavelength / smaller wavelength [1]
- (ii) Hydrogen since energy is proportional to frequency / greater frequency / E = hf [1]
- (d) In Ne greater shielding of *outer* electron (1) outweighs larger nuclear charge (1) / He has greater effective nuclear charge (1) / He *outer* electron closer to nucleus (1)
 - max 1 if no reference to *outer* electron [2]
 (Maximum 2 marks)
- (e) (i) ²¹⁸Po [1]
- (ii) Since radon is a gas / inhaled, α particles will be given off in the lungs (which may cause cancer) [1]

Total [12]

9. (a) Low temperature (1)
 As temperature is decreased equilibrium moves in exothermic direction. (1)
- High pressure (1)
 As pressure is increased equilibrium moves towards side with smaller number of gas moles (1) [4]
- QWC The information is organised clearly and coherently, using specialist vocabulary where appropriate [1]
- (b) $\Delta H_{\text{reaction}} = \Delta H_{\text{f products}} - \Delta H_{\text{f reactants}}$ (1)
 $-46 = \Delta H_{\text{f ethanol}} - (52.3 - 242)$
 $\Delta H_{\text{f ethanol}} = -46 - 189.7$ (1)
 $\Delta H_{\text{f ethanol}} = -235.7 \text{ kJ mol}^{-1}$ (1) [3]
- (c) Bonds broken = $1648 + 612 + 926 = 3186 \text{ kJ mol}^{-1}$ (1)
 Bonds formed = $2060 + 348 + 360 + 463 = 3231 \text{ kJ mol}^{-1}$ (1)
 $\Delta H_{\text{reaction}} = 3186 - 3231 = -45 \text{ kJ mol}^{-1}$ (1) [3]
- (d) (i) Average bond enthalpies used (not actual ones) [1]
 (ii) Yes, since answers are close to each other [1]
- (e) Catalyst is in different (physical) state to reactants [1]



Total [16]

10. (a) Weighing bottle would not have been washed / difficult to dissolve solid in volumetric flask / final volume would not necessarily be 250 cm³ [1]
- (b) Pipette [1]
- (c) To show the end point / when to stop adding acid / when it's neutralised [1]
- (d) So that a certain volume of acid can be added quickly before adding drop by drop / to save time before doing accurate titrations / to give a rough idea of the end point [1]
- (e) To obtain a more reliable value [1]
- (f) (i) Moles = $0.730/36.5 = 0.0200$ (1)
- Concentration = $0.02/0.1 = 0.200 \text{ mol dm}^{-3}$ (1) [2]
- (ii) Moles = $0.2 \times 0.0238 = 0.00476$ [1]
- (iii) 0.00476 [1]
- (iv) $0.00476 \times 10 = 0.0476$ [1]
- (v) $M_r = 1.14/0.0476 = 23.95$ [1]
- (vi) Lithium [1]

- mark consequentially throughout (f)

Total [12]

Section B Total [70]