



# **GCE MARKING SCHEME**

**CHEMISTRY  
AS/Advanced**

**JANUARY 2013**

**GCE CHEMISTRY - CH1**  
**JANUARY 2013 MARK SCHEME**

**SECTION A**

Q.1 39 [1]

Q.2 C [1]

Q.3  $A_r = \frac{(12.0 \times 6) + (88.0 \times 7)}{100} (1) = \frac{72.0 + 616.0}{100} = 6.88 (1)$  [2]

Q.4 (a)  $\Delta H = \Delta H_2 + \Delta H_3 - \Delta H_1$  [1]

(b)  $\frac{1}{2}\text{N}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightarrow \text{NO}(\text{g})$  state symbols requires [1]

Q.5 The position of equilibrium moves to the right / more COS is formed (1)  
(By Le Chatelier's principle) the system 'removes' added 'material' to restore the  
position of equilibrium / accept explanation in terms of pressure (1) [2]

Q.6 Ti  $\frac{60}{48}$  O  $\frac{40}{16}$  (1)  
  
= 1.25 = 2.5  $\therefore 1 : 2$   
  
 $\therefore \text{TiO}_2$  (1) [2]

**Section A Total [10]**

**SECTION B**

Q.7 (a) (i) A helium (atom) nucleus / 2 protons and 2 neutrons /  ${}^4\text{He}^{2+}$  [1]

(ii) b.....22 (1) X.....Ne (1) [2]

(iii)  $(4 \times 2.6) = 10.4$  [1]

(b) The frequency of the green line at 569 nm is HIGHER. than the frequency of the yellow-orange line at 589 nm. Another line is seen at 424 nm, this is caused by an electronic transition of HIGHER. energy than the line at 569 nm. [1]

(c) (i) 
$$\begin{array}{ccccccc} \text{Na}_2\text{CO}_3 & & \text{NaHCO}_3 & & 2\text{H}_2\text{O} & & \\ \downarrow & & \downarrow & & \downarrow & & \\ 106 & + & 84 & + & 36 & (1) & \rightarrow 226 \end{array}$$
 [1]

(or by other appropriate method – note mark is for the working)

(ii) Atom economy =  $\frac{\text{'M}_r \text{ required product} \times 100}{\text{Total 'M}_r \text{ of the reactants}}$  (1)  
 $= \frac{318 \times 100}{452} = 70.4 / 70.35 (\%)$  (1) [2]

(iii) Carbon dioxide is produced (and released into the air) and this contributes to the greenhouse effect / increases acidity of sea (1)  
 It should be trapped / a use found for it. (1) [2]

(d) (i) Water is acting as a proton donor (1) and this combines with the carbonate ion /  $\text{CO}_3^{2-}$ , giving the hydrogencarbonate ion /  $\text{HCO}_3^-$  (1) [2]

(ii) The pH scale runs from 0-14 / measure of acidity / alkalinity (1)  
 pH <7 acid / >7 alkali (1)  
 acid stronger as pH value decreases / alkali stronger as pH value increases / 11.4 is strong alkali (1) [3]

**Total [15]**

Q.8 (a) (i) He may have lost carbon dioxide through leaks, this would have given a lower volume than expected. (1)  
He used lower concentration of acid / diluted the acid with water and the rate of carbon dioxide evolution was slower than expected. (1) [2]

(ii) The concentration of acid is higher in the first half (1) the collision rate is higher (1) [2]

(iii) eg  $k = \frac{V}{T}$  (1)  $\therefore k = \frac{130}{298}$  / 0.436

$\therefore V = 0.436 \times 323 = 141 \text{ (cm}^3\text{)}$  (1)

or  $\frac{V_1}{V_2} = \frac{T_1}{T_2}$  (1)  $\therefore V_1 = \frac{323 \times 130}{298} = 141 \text{ (cm}^3\text{)}$  (1) [2]

(b) (i) 260 (cm<sup>3</sup>) [1]

(ii) 0.45 (g) (0.43–0.48) [1]

(c) The diagram shows two reasonable distribution curves with T<sub>2</sub> flatter and 'more to the right' than T<sub>1</sub>. (1)  
Activation energy correctly labelled, or mentioned in the writing (1)  
Fraction of molecules having the required activation energy is much greater at a higher temperature (thus increasing the frequency of successful collisions) (in words) (1) [3]

*The candidate has selected a form and style of writing that is appropriate to purpose and complexity of the subject matter QWC* [1]

(d) Place the mixture on a balance and measure the (loss in) mass (1)  
at appropriate time intervals (1)

OR BY OTHER SUITABLE METHOD

eg. sample at intervals / quench (1) titration (1) [2]


**Total [14]**

Q.9 (a) (i) They are both elements in their standard states. [1]

(ii)  $\Delta H = \sum \Delta H_f \text{ products} - \sum \Delta H_f \text{ reactants}$  (1)  
 $= (-286 + 0) - (-368 + 0)$   
 $= -286 + 368 = (+)82 \text{ (kJ mol}^{-1}\text{)}$  (1) [2]

or by a cycle where correct cycle drawn (1) correct answer (1)

(b) (i)



exothermic profile drawn (1)  
uncatalysed / catalysed line labelled (1) [2]

(ii) I number of moles of benzene = 2000 [1]

II mole ratio is 1 : 1 (1)

$$\therefore \text{moles of phenol produced} = \frac{2000 \times 95}{100} = 1900 \text{ (1)}$$

$$\text{mass} = M_r \times \text{number of moles} = 94 \times 1900 = 178.6 / 179 \text{ kg (1)}$$

*alternatively*

78 (g / kg) of benzene gives 94 (g / kg) of phenol (1)

$\therefore$  1 (g / kg) of benzene gives  $94/78$  (g / kg) of phenol

$\therefore$  156 (kg) of benzene gives  $94 \times 156/78$  (kg) of phenol = 188 (kg) (1)

but 95% yield  $\therefore \frac{188 \times 95}{100} = 178.6 / 179$  (kg) (1) [3]

(iii) Look for at least four relevant positive points [4]

- e.g.
- the process uses a (heterogeneous) catalyst, which can easily be separated from the gaseous products (thus saving energy)
  - the only other product of the reaction is gaseous nitrogen, which is non-toxic / safe / not a harmful product
  - the process uses nitrogen(I) oxide which is used up, rather than being released into the atmosphere from the other process (and causing global warming)
  - the process is exothermic and the heat produced can be used elsewhere
  - a relatively moderate operating temperature reduces overall costs
  - high atom economy

*Legibility of text; accuracy of spelling, punctuation and grammar;  
clarity of meaning* QWC [1]

**Total [14]**

Q.10 (a)  $K \rightarrow 1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$  (1)  
There is one outer electron and the loss of this electron gives a stable potassium ion with a full outer shell/ ion more stable than the atom (1) [2]

(b) (i)  $\Delta T = 4.8 \text{ }^\circ\text{C}$  (1)

$$\Delta H = - \frac{250 \times 4.2 \times 4.8}{0.125} = - 40320 \text{ J mol}^{-1} / - 40.3 \text{ kJ mol}^{-1} \text{ (2) [3]}$$

✓ for negative sign

✓ correct value with relevant units

(ii) e.g. The volume used was not precise in measurement as the readings on a beaker are only approximate (1)  
The experiment was performed in a beaker and this was not insulated and heat was lost to the surroundings (1) [2]

there may be other acceptable answers here, for example based on slow dissolving

(c) (i) 0.050 [1]

(ii)  $(0.050 \times 24.0) = 1.20 \text{ (dm}^3\text{)}$  [1]

(iii)  $\% \text{ v/v} = \frac{1.20 \times 0.001 \times 100}{2} \text{ (1)} = 0.06 \text{ (1)} \text{ [2]}$

(d) An increase in the concentration of (aqueous) carbon dioxide causes the position of equilibrium to move to the right. (1)  
This causes calcium carbonate to become aqueous calcium (and hydrogencarbonate) ions / dissolve (1)  
weakening shells / causing difficulty in formation of shells (1) [3]

*Organisation of information clearly and coherently; using specialist vocabulary where appropriate* QWC [1]

**Total [15]**

- Q.11 (a) (i) I burette / (graduated) pipette [1]  
 II volumetric / graduated / standard flask [1]
- (ii) 0.0064 [1]
- (iii) 1.20 g / 100 cm<sup>3</sup> solution [1]
- (iv) 12.0 g / 100 cm<sup>3</sup> solution [1]
- (b) (i) The catalyst is in a different physical state to the reactants. [1]
- (ii) Bonds broken    2 H-H → 872            1 C-O → 360  
                           1 C-H → 412            1 O-H → 463  
                           1 C=O → 743
- Total +2850 kJ (1)
- Bonds made    3 C-H → 1236  
                           1 C-O → 360  
                           3 O-H → 1389
- Total -2985 kJ (1)
- $\Delta H = 2850 - 2985 = -135 \text{ kJ mol}^{-1}$  (1) [3]
- (c) Relative molecular mass is a relative quantity (based on  $1/12$ th of the  $^{12}\text{C}$  atom as one unit). [1]
- (d) (i) The rate of the forward reaction is equal to the rate of the backward reaction. [1]
- (ii) C<sub>2</sub>H<sub>4</sub>O [1]

**Total [12]**

**Total Section B [70]**