



GCE MARKING SCHEME


**CHEMISTRY
AS/Advanced**

SUMMER 2011

CHEMISTRY - CH2


SECTION A

Q.1	(a)	Calcium carbonate	[1]
	(b)	Sodium carbonate	[1]
Q.2		Metallic (1) Covalent and van der Waals (1)	[2]
Q.3		$\text{Ca}_3(\text{PO}_4)_2$	[1]
Q.4		D	[1]
Q.5		Materials that change their properties in response to a change in conditions / environment / surroundings	[1]
Q.6	(a)	Alkene / double bond (1) Alcohol / hydroxyl / hydroxy (1)	[2]
	(b)	$\text{C}_5\text{H}_{10}\text{O}$	[1]
			Total [10]





UV light (1)

any of following for 4 max




Free radical substitution / photochlorination (1)



[6]

QWC: Selection of form and style of writing appropriate to purpose and to complexity of subject matter.

[1]



Nucleophilic substitution / hydrolysis

[2]

(c) Heat with NaOH (1)

Add HNO_3 then AgNO_3 (1)

White precipitate seen (1)


[3]

(d) Ozone layer depleted / (leads to) increased incidence of skin cancer


Contributes to greenhouse effect / increases global warming

[1]

Total [13]

- Q.9** (a) C=O absorption at 1650–1750 cm⁻¹
 C–O absorption at 1000–1300 cm⁻¹
 O–H absorption at 2500–3500 cm⁻¹
 3 correct peaks labelled [2]
 (2 correct peaks labelled 1 mark)
- (b) Molecular ion at m/z 60 shows that M_r is 60 (1)
 Peak at m/z 15 shows CH₃ group / peak at m/z 45 shows COOH group (1) [2]
- (c) (i)  [1]
 (Accept 1 hydrogen bond)
- (ii) (Intermolecular bond formed) when hydrogen attached to a highly electronegative atom (oxygen) (1)
 is bonded to an electronegative atom in another molecule (1)
 forming very strong dipole – dipole attraction (1) [3]
QWC: Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning [1]
- (d) (i) Acidified and heat / reflux [1]
 (ii) Colour change from orange to green [1]
- (e) Propane would be lower as it cannot form hydrogen bonds / only forms van der Waals forces between molecules (1)
 Butan-1-ol would be higher as it (also has hydrogen bonds but) has more van der Waals forces between molecules (1) [2]

Total [13]



(ii)

<i>Element</i>	<i>Initial Oxidation State</i>	<i>Final Oxidation State</i>
Nitrogen	-3	2
Hydrogen	1	1
Oxygen	0	-2

All three rows correct (2)
(1 mark if two rows correct)


Nitrogen oxidised as its oxidation state has increased (1) [3]

(iii) NH_3 has 3 bonding and 1 non bonding pair of electrons (1)

BF_3 has 3 bonding pairs only (1)

Electron pairs position themselves as far apart as possible
(to minimise repulsion) (1) [3]

(b) (i) A covalent bond where one of the atoms has donated both electrons
in the shared pair [1]



(iii) Tetrahedral (1) [2]

$109\frac{1}{2}^\circ$ (1) (accept 109°) [2]

(iv) Water is polar / a polar solvent (1)

Anion is attracted to H^{\oplus} / cation is attracted to O^{\ominus} (1) [2]

Total [14]

- Q.11** (a) (i) Lilac flame (1)
 White solid / white fumes / potassium melts (1) [2]
- (ii) $4\text{K} + \text{O}_2 \longrightarrow 2\text{K}_2\text{O}$ [1]
- (iii) More reactive (1)
 Electrons in rubidium lost more easily / ionisation energy is less /
 explanation e.g. increased sheilding (1) [2]
 (Need reason to get first mark but accept more reactive as reactivity
 increases down group for 1 mark)
- (b) (i) No. moles = $\frac{0.098}{23} = 0.00426$ [1]
- (ii) Moles $\text{H}_2 = 0.00213$ (1)
 Volume $\text{H}_2 = 0.00213 \times 24 = 0.0511 \text{ dm}^3$ (1) [2]
- (iii) Moles $\text{NaOH} = 0.00426$ (1)
 Concentration $\text{NaOH} = \frac{0.00426}{0.200} = 0.0213 \text{ mol dm}^{-3}$ (1) [2]
- (c) (i) Do the experiment in a fume cupboard [1]
- (ii) I 6:6 [1]
 II Electrostatic forces between the oppositely charged ions (1)
 ionic bonds are / ionic lattice is very strong so large amount of
 energy needed (1) [2]

Total [14]

Section B Total [70]