

# GCE Examinations

# Further Pure Mathematics Module FP2

Advanced Subsidiary / Advanced Level

Paper D

Time: 1 hour 30 minutes

## *Instructions and Information*

---

Candidates may use any calculator except those with a facility for symbolic algebra and/or calculus.

Full marks may be obtained for answers to ALL questions.

Mathematical and statistical formulae and tables are available.

This paper has 7 questions.

## *Advice to Candidates*

---

You must show sufficient working to make your methods clear to an examiner.  
Answers without working will gain no credit.



*Written by Rosemary Smith & Shaun Armstrong*

© Solomon Press

*These sheets may be copied for use solely by the purchaser's institute.*

1. 
$$y = \frac{\operatorname{cosech} x}{x^2 + 1}.$$

(a) Find  $\frac{dy}{dx}$ . **(4 marks)**

(b) Find the value of  $\frac{dy}{dx}$  when  $x = 0.5$ , giving your answer to 2 decimal places. **(1 mark)**

---

2. A curve has intrinsic coordinates  $(s, \psi)$  and radius of curvature  $\rho$ .

Given that  $\rho = 2(s + a)$ , where  $a$  is constant, show that the intrinsic equation of the curve can be written in the form

$$s = Ae^{2\psi} - a,$$

where  $A$  is constant. **(5 marks)**

---

3. (a) Prove that

$$\sinh 3x \equiv 4 \sinh^3 x + 3 \sinh x. \quad \textbf{(5 marks)}$$

(b) Hence, or otherwise, solve the equation

$$\sinh 3x = 7 \sinh^2 x,$$

giving your answers in terms of natural logarithms where appropriate. **(6 marks)**

---

4. (a) Find  $\int \frac{1}{\sqrt{9-4x^2}} dx$ . **(3 marks)**

(b) Find  $\int \frac{1-2x}{\sqrt{9-4x^2}} dx$ . **(3 marks)**

(c) Hence, or otherwise, solve the differential equation

$$\sqrt{9-4x^2} \frac{dy}{dx} = y(1-2x),$$

given that  $y = 1$  when  $x = 0$ . **(6 marks)**

---

5. The curve  $C$  has equation  $y^2 = 4ax$ , where  $a$  is a positive constant.

(a) Show that an equation of the tangent to  $C$  at the point  $P(ap^2, 2ap)$ ,  $p \neq 0$ , is

$$yp = x + ap^2. \quad (4 \text{ marks})$$

The point  $Q(aq^2, 2aq)$ , is on  $C$  where  $q \neq 0$  and  $p \neq q$ . The chord  $PQ$  passes through the focus of  $C$ .

Show that

(b)  $pq = -1$ , (5 marks)

(c) the tangent to  $C$  at  $P$  and the tangent to  $C$  at  $Q$  meet on the directrix of  $C$ . (4 marks)

---

6. 
$$I_n = \int_0^{\frac{\pi}{4}} \sec^n x \, dx, \quad n \geq 0.$$

(a) Show that

$$(n-1)I_n = (\sqrt{2})^{n-2} + (n-2)I_{n-2}, \quad n \geq 2. \quad (7 \text{ marks})$$

(b) Hence find the exact value of  $I_3$ , giving your answer in terms of natural logarithms.

(6 marks)

---

7. (a) Show that

$$\int \sqrt{a^2 + x^2} \, dx = \frac{x}{2} \sqrt{a^2 + x^2} + \frac{a^2}{2} \operatorname{arsinh}\left(\frac{x}{a}\right) + c. \quad (9 \text{ marks})$$

The parametric equations of the curve  $C$  are

$$x = 2t, \quad y = t^2, \quad 0 \leq t \leq 3.$$

(b) Show that the length of  $C$  is given by

$$2 \int_0^3 \sqrt{1+t^2} \, dt. \quad (4 \text{ marks})$$

(c) Find the length of  $C$ . (3 marks)

---

**END**