

EDEXCEL FOUNDATION

Stewart House 32 Russell Square London WC1B 5DN

January 2002

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6679

* indicates printed answer

Paper No. M3

Question number	Scheme	Marks
1.	$0.2a = \frac{5}{x+1}$ $0.2v \frac{dv}{dx} = \frac{5}{x+1}$ $\int v dv = \int \frac{25}{x+1} dx$ $\frac{1}{2} v^2 = 25 \ln(x+1) (+ C)$ $x=0, v=5 \Rightarrow C = 12.5$ $\frac{225}{2} = 25 \ln(x+1) + 12.5$ $x = 53.6 \text{ (3sf)}$	<p style="text-align: center;">M1</p> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">}</div> <div style="margin-right: 5px;">→</div> <div>M1</div> </div> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">}</div> <div style="margin-right: 5px;">→</div> <div>M1</div> </div> <p style="text-align: center;">A1 A1</p> <p style="text-align: center;">A1</p> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">}</div> <div style="margin-right: 5px;">→</div> <div>M1</div> </div> <p style="text-align: center;">A1</p> <p style="text-align: right;">(8)</p>
2. (a)	<p>PE Loss = $0.5g(2+x)$; $EPE_{atc} = \frac{19.6x^2}{4}$</p> $0.5g(2+x) = \frac{19.6x^2}{4}$ $k(x^2 - 2 - 2) = 0$ <p style="text-align: center;">solving</p> $AC = 4m$	<p style="text-align: center;">B1; B1</p> <p style="text-align: center;">M1</p> <p style="text-align: center;">A1</p> <p style="text-align: center;">M1</p> <p style="text-align: center;">A1 ✓</p> <p style="text-align: right;">(6)</p>
(b)	$T_c = \frac{19.6 \times 2}{2} = 19.6$ $19.6 - 0.5g = 0.5a$ $a = 29.4 \text{ ms}^{-2}$	<p style="text-align: center;">B1 ✓</p> <p style="text-align: center;">M1</p> <p style="text-align: center;">A1</p> <p style="text-align: right;">(3)</p> <p style="text-align: right;">(9)</p>
3. (a)	Line of action of weight must pass through c which is not above centre of rod (or equivalent)	<p style="text-align: center;">B1</p> <p style="text-align: right;">(1)</p>
(b)	<p><u>Method A:</u></p> <p>R (along ac): $T_1 = 2mg \sin \alpha = \frac{6mg}{5}$</p> <p>R (along bc): $T_2 = 2mg \cos \alpha = \frac{8mg}{5}$*</p> <p>[Equiv. to moments about A, B respectively]</p> <p>or <u>Method B:</u> R(A), $T_1 \sin \alpha + T_2 \cos \alpha = 2mg$</p> <p>L(→), $T_1 \cos \alpha = T_2 \sin \alpha$</p> <p>solving to find T_1 or T_2</p> $T_1 = \frac{6mg}{5}; T_2 = \frac{8mg}{5}$ *	<p style="text-align: center;">M1 M1 A1</p> <p style="text-align: center;">M1 A1</p> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">}</div> <div style="margin-right: 5px;">→</div> <div>M1</div> </div> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">}</div> <div style="margin-right: 5px;">→</div> <div>M1</div> </div> <p style="text-align: center;">A1; A1</p> <p style="text-align: right;">(5)</p>
(c)	$\frac{8mg}{5} = \frac{k \cos(BC-a)}{a}$ $BC = 2a \sin \alpha$ $k = 8$	<p style="text-align: center;">M1 A1</p> <p style="text-align: center;">B1</p> <p style="text-align: center;">A1</p> <p style="text-align: right;">(4)</p> <p style="text-align: right;">(10)</p>

EDEXCEL FOUNDATION

Stewart House 32 Russell Square London WC1B 5DN



January 2002

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6679

Paper No. M3

Question number	Scheme	Marks
4.(a)	$\int_0^r (\pi) y^2 x dx = \pi \int_0^r (\pi) y^2 dx$ $\int_0^r r x^2 dx = \pi \int_0^r r x dx$ $\left[(r) \frac{x^3}{3} \right]_0^r = \pi \left[(r) \frac{x^2}{2} \right]_0^r$ $\bar{x} = \frac{2r}{3} *$	<p>M1 A1 M1 A1 A1 A1 (6)</p>
(b)	 <p>vertical thro' CM and lowest point of plane face</p> $\tan \alpha = \frac{r}{r/3}$ $\alpha = 72^\circ \text{ (nearest degree)}$	<p>M1 M1 A1 A1 (4)</p>
5.	 <p> $R(1), R \sin 25^\circ - F \sin 25^\circ = mg$ $R(\leftarrow), R \cos 25^\circ + F \cos 25^\circ = \frac{mv^2}{40}$ $F = 0.6R$ used Eliminating R Solving for v $v = 24.1 \text{ ms}^{-1}, 24 \text{ ms}^{-1}$ </p>	<p>M1 A2 M1 A2 M1 M1 M1 A1 (10)</p>
6.(a)	<p>If SHM, $a = 1.2$</p> <p>Using $v^2 = \omega^2(a^2 - x^2)$</p> $0.27 = \omega^2(1.2^2 - 0.6^2) \text{ or } 0.2 = \omega^2(1.2^2 - 0.8^2)$ <p>Solve for $\omega (= 0.5)$ and use in other eqn²</p> <p>Shown to be correct</p> <p>(b) $v = a\omega = 1.2 \times 0.5 = 0.6 *$</p> <p>(c) $\ddot{x} = \omega^2 \times 0.6 = 0.15 \text{ m s}^{-2}$</p> <p>(d) $0.6 = a \sin \omega t$ or $0.8 = a \sin \omega t$</p> $t = \frac{1}{\omega} \left(\sin^{-1} \frac{0.8}{a} - \sin^{-1} \frac{0.6}{a} \right)$ $= 0.412 \text{ s (3SF)}$	<p>B1 M1 A1 M1 A1 c.s.o. (5)</p> <p>M1 A1 (2)</p> <p>M1 A1 V (2)</p> <p>M1 M1 A1 V A1 (4)</p> <p>(13)</p>

EDEXCEL FOUNDATION

Stewart House 32 Russell Square London WC1B 5DN

January 2002

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6679

Paper No. M3

Question number	Scheme	Marks
7.(a)	$\frac{1}{2} m \frac{7as}{2} - \frac{1}{2} m v^2 = mga$ $\leftarrow, R = \frac{mv^2}{a} = \frac{3mg}{-2}$	<p>M1 A1 M1 A1 (4)</p>
(b)	$\frac{1}{2} m \frac{7as}{2} - \frac{1}{2} m v^2 = mga (1 + \cos \theta)$ $\leftarrow, mg \cos \theta = \frac{mv^2}{a}$ <p>Eliminating v^2</p> <p>Solving to give $\cos \theta = k, \theta = 60^\circ *$</p>	<p>M1 A1 M1 A1 M1 M1 A1 (7)</p>
(c)	$v \cos 60^\circ t = a \sin 60^\circ$ $v^2 = ag \cos 60^\circ$ <p>Making t explicit</p> $t = \sqrt{\frac{6a}{g}}$	<p>M1 B1 M1 A1 (4)</p> <p>(15)</p>