

# GCE Examinations

## Mechanics

### Module M3

Advanced Subsidiary / Advanced Level

Paper A

Time: 1 hour 30 minutes

#### *Instructions and Information*

---

Candidates may use any calculator except those with a facility for symbolic algebra and/or calculus.

Full marks may be obtained for answers to ALL questions.

Mathematical and statistical formulae and tables are available.

This paper has 7 questions.

When a numerical value of  $g$  is required, use  $g = 9.8 \text{ m s}^{-2}$ .

#### *Advice to Candidates*

---

You must show sufficient working to make your methods clear to an examiner. Answers without working will gain no credit.



*Written by Shaun Armstrong & Chris Huffer*

© Solomon Press

*These sheets may be copied for use solely by the purchaser's institute.*

1.




Fig. 1

A particle of mass 0.6 kg is attached to one end of a light elastic spring of natural length 1 m and modulus of elasticity 30 N. The other end of the spring is fixed to a point  $O$  which lies on a smooth plane inclined at an angle  $\alpha$  to the horizontal where  $\tan \alpha = \frac{3}{4}$  as shown in Figure 1.

The particle is held at rest on the slope at a point 1.2 m from  $O$  down the line of greatest slope of the plane.

(a) Find the tension in the spring. **(2 marks)**

(b) Find the initial acceleration of the particle. **(5 marks)**

---

2. A particle  $P$  of mass 0.5 kg moves along the positive  $x$ -axis under the action of a single force directed away from the origin  $O$ . When  $P$  is  $x$  metres from  $O$ , the magnitude of the force is  $3x^{\frac{1}{2}}$  N and  $P$  has a speed of  $v$   $\text{m s}^{-1}$ .

Given that when  $x = 1$ ,  $P$  is moving away from  $O$  with speed  $2$   $\text{m s}^{-1}$ ,

(a) find an expression for  $v^2$  in terms of  $x$ , **(5 marks)**

(b) show that when  $x = 4$ ,  $P$  has a speed of  $7.7$   $\text{m s}^{-1}$ , correct to 1 decimal place. **(2 marks)**

---

3. A particle is performing simple harmonic motion along a straight line between the points  $A$  and  $B$  where  $AB = 8$  m. The period of the motion is 12 seconds.

(a) Find the maximum speed of the particle in terms of  $\pi$ . **(4 marks)**

The points  $P$  and  $Q$  are on the line  $AB$  at distances of 3m and 6m respectively from  $A$ .

(b) Find, correct to 3 significant figures, the time it takes for the particle to travel directly from  $P$  to  $Q$ .

**(6 marks)**

---

4. Whilst in free-fall a parachutist falls vertically such that his velocity,  $v \text{ m s}^{-1}$ , when he is  $x$  metres below his initial position is given by

$$v^2 = kg(1 - e^{-\frac{2x}{k}}),$$

where  $k$  is a constant.

Given that he experiences an acceleration of  $f \text{ m s}^{-2}$ ,

- (a) show that  $f = ge^{-\frac{2x}{k}}$ . (4 marks)

After falling a large distance, his velocity is constant at  $49 \text{ m s}^{-1}$ .

- (b) Find the value of  $k$ . (3 marks)

- (c) Hence, express  $f$  in the form  $(\lambda - \mu v^2)$  where  $\lambda$  and  $\mu$  are constants which you should find. (4 marks)

5.




Fig. 2

A firework is modelled as a uniform solid formed by joining the plane surface of a right circular cone of height  $2r$  and base radius  $r$ , to one of the plane surfaces of a cylinder of height  $h$  and base radius  $r$  as shown in Figure 2.

Using this model,

- (a) show that the distance of the centre of mass of the firework from its plane base is

$$\frac{3h^2 + 4hr + 2r^2}{2(3h + 2r)}. \quad (9 \text{ marks})$$

The firework is to be launched from rough ground inclined at an angle  $\alpha$  to the horizontal. Given that the firework does not slip or topple and that  $h = 4r$ ,

- (b) Find, correct to the nearest degree, the maximum value of  $\alpha$ . (4 marks)

*Turn over*

6.




Fig. 3

The two ends of a light inextensible string of length  $3a$  are attached to fixed points  $Q$  and  $R$  which are a distance of  $a\sqrt{3}$  apart with  $R$  vertically below  $Q$ . A particle  $P$  of mass  $m$  is attached to the string at a distance of  $2a$  from  $Q$ .

$P$  is given a horizontal speed,  $u$ , such that it moves in a horizontal circle with both sections of the string taut as shown in Figure 3.

- (a) Show that  $\angle PRQ$  is a right angle. (2 marks)
- (b) Find  $\angle PQR$  in degrees. (1 mark)
- (c) Find, in terms of  $a$ ,  $g$ ,  $m$  and  $u$ , the tension in the section of string
- (i)  $PQ$ ,
- (ii)  $PR$ . (7 marks)
- (d) Show that  $u^2 \geq \frac{ga}{\sqrt{3}}$ . (3 marks)

7. A particle of mass  $2$  kg is attached to one end of a light elastic string of natural length  $1$  m and modulus of elasticity  $50$  N. The other end of the string is attached to a fixed point  $O$  on a rough horizontal plane and the coefficient of friction between the particle and the plane is  $\frac{10}{49}$ .

The particle is projected from  $O$  along the plane with an initial speed of  $5 \text{ m s}^{-1}$ .

- (a) Show that the greatest distance from  $O$  which the particle reaches is  $1.84$  m. (9 marks)
- (b) Find, correct to 2 significant figures, the speed at which the particle returns to  $O$ . (5 marks)

END