Centre No.						Pape	er Refer	ence			Surname	Initial(s)
Candidate No.	;			6	6	8	0	/	0	1	Signature	

Paper Reference(s)

6680/01

Edexcel GCE

Mechanics M4

Advanced/Advanced Subsidiary

Thursday 12 June 2008 – Morning

Time: 1 hour 30 minutes

Materials required f	for	examination
----------------------	-----	-------------

mathematical formulae stored in them.

Mathematical Formulae (Green)

Items included with question papers

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions. Write your answers in the spaces provided in this question paper. Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 7 questions in this question paper. The total mark for this paper is 75.

There are 24 pages in this question paper. Any blank pages are indicated.


Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.

H30094RA W850/R6680/57570 3/3/3/3/3/3/3





Examiner's use only

Team Leader's use only

1

3

4

5

6



	A ship P is moving with velocity $(5\mathbf{i} - 4\mathbf{j})$ km h ⁻¹ and a ship Q is moving with velocity $(3\mathbf{i} + 7\mathbf{j})$ km h ⁻¹ . Find the direction that ship Q appears to be moving in, to an observer on ship P , giving your answer as a bearing.
	(5)
_	

Question 1 continued	Le bla
zuestion i continueu	
	- 1



uestion 2 continued	Lea blai
	Q2

		Lea blaı
3.	At time $t = 0$, a particle of mass m is projected vertically downwards with speed U from a point above the ground. At time t the speed of the particle is v and the magnitude of the air resistance is modelled as being mkv , where k is a constant.	Diai
	Given that $U < \frac{g}{2k}$, find, in terms of k, U and g, the time taken for the particle to double	
	its speed.	
	(8)	

Question 3 continued	Leave blank
	Q3
(Total 8 marks)	

Leave blank

4.




Figure 1

A small smooth ball B, moving on a horizontal plane, collides with a fixed vertical wall. Immediately before the collision the angle between the direction of motion of B and the wall is 2θ , where $0^{\circ} < \theta < 45^{\circ}$. Immediately after the collision the angle between the direction of motion of B and the wall is θ , as shown in Figure 1. Given that the coefficient of restitution between B and the wall is $\frac{3}{8}$, find the value of $\tan \theta$.

(8)

Question 4 continued	Leave blank
	Q4
/TE / 10 1 1	
(Total 8 marks)	

5.	A light elastic spring has natural length l and modulus of elasticity mg . One end of a spring is fixed to a point O on a rough horizontal table. The other end is attached to a partial P of mass m which is at rest on the table with $OP = l$. At time $t = 0$ the particle is project with speed $\sqrt{(gl)}$ along the table in the direction OP . At time t the displacement of P its initial position is x and its speed is y . The motion of P is subject to air resistance	cle ted om
	magnitude $2mv\omega$, where $\omega = \sqrt{\frac{g}{l}}$. The coefficient of friction between P and the table	
	0.5. V <i>l</i>	
	(a) Show that, until <i>P</i> first comes to rest,	
	$\frac{\mathrm{d}^2 x}{\mathrm{d}t^2} + 2\omega \frac{\mathrm{d}x}{\mathrm{d}t} + \omega^2 x = -0.5g.$	(6)
	(b) Find x in terms of t , l and ω .	
		(6)
	(c) Hence find, in terms of ω , the time taken for <i>P</i> to first come to instantaneous rest.	
		(3)
		_
		_
		_
		_
		_
		_
		_
		_

Question 5 continued	Leave blank

Question 5 continued	b]

Question 5 continued	Leav blan
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	-
	-
	_
	- _
	_ Q:

6.




Figure 2

A river is 30 m wide and flows between two straight parallel banks. At each point of the river, the direction of flow is parallel to the banks. At time t = 0, a boat leaves a point O on one bank and moves in a straight line across the river to a point P on the opposite bank. Its path OP is perpendicular to both banks and OP = 30 m, as shown in Figure 2. The speed of flow of the river, r m s⁻¹, at a point on OP which is at a distance x m from O, is modelled as

$$r = \frac{1}{10}x, \quad 0 \le x \le 30.$$

The speed of the boat relative to the water is constant at 5 m s⁻¹. At time t seconds the boat is at a distance x m from O and is moving with speed v m s⁻¹ in the direction OP.

(a) Show that

$$100v^2 = 2500 - x^2. ag{3}$$

(b) Hence show that

$$\frac{d^2x}{dt^2} + \frac{x}{100} = 0. {4}$$

(c) Find the total time taken for the boat to cross the river from O to P.

(9)

Leave blank



Question 6 continued	Leave blank

Question 6 continued	

Question 6 continued	Leave blank
	06
(Total 16 marks)	Q6

7.




Figure 3

A uniform rod AB, of length 2a and mass kM where k is a constant, is free to rotate in a vertical plane about the fixed point A. One end of a light inextensible string of length 6a is attached to the end B of the rod and passes over a small smooth pulley which is fixed at the point P. The line AP is horizontal and of length 2a. The other end of the string is attached to a particle of mass M which hangs vertically below the point P, as shown in Figure 3. The angle PAB is 2θ , where $0^{\circ} \le \theta \le 180^{\circ}$.

(a) Show that the potential energy of the system is

$$Mga(4\sin\theta - k\sin 2\theta) + \text{constant.}$$
 (5)

The system has a position of equilibrium when $\cos \theta = \frac{3}{4}$.

(b) Find the value of k.

(5)

Leave blank

(c) Hence find the value of $\cos\theta$ at the other position of equilibrium.

(3)

(d) Determine the stability of each of the two positions of equilibrium.

(5)

Question 7 continued	Leave blank

	Lea
Question 7 continued	

20

	Leave blank
Question 7 continued	

	estion 7 continued	
(T) 110 110		
(Total 18 marks)		
TOTAL FOR PAPER: 75 MARKS		 RKS

