


PH2

Question		Marking details	Marks Available	
1	(a)	(i)	0.04[0 m]	1
		(ii)	$T = 0.20$ s [or by impl.] (1) $f = 5.0$ (1) Hz (1) (e.c.f. on T)	3
	(b)	If peak arriving at 0.050 s at B is the peak that passed A at 0.00 s [or equiv] (1), $v = \frac{0.30 \text{ m}}{0.050 \text{ s}}$ [free-standing](1) [Accept: B could be $\lambda/4$ from A , so $\lambda = 1.2$ m (1); $v = f\lambda = 5.0 \times 1.2 \text{ m s}^{-1}$ (1).]	2	
	(c)	(i)	Distance [along the direction of wave propagation] between two [consecutive] point (1) oscillating in phase (1) [“Distance between two peaks / troughs \rightarrow 1]	2
		(ii)	$\lambda = 1.2$ m (e.c.f. on f)	1
			[9]	

Question		Marking details	Marks Available
2	(a)	(i) Wavefronts [or waves] from each slit spread out (1) [accept: waves diffract at each slit]and overlap (1) [or superpose or interfere].	2
		(ii) I. Sources which emit waves, which are at the same point in their cycle at the same time [accept: "emit peaks at the same time"] II. A maximum on central axis or microwave source central w.r.t. S_1 and S_2 .	1 1
		(iii) Correct insertion of values into $\lambda = \frac{ay}{D}$ (1) [or by implication] $\lambda = 0.012$ m (1)	2
		(iv) I. Constructive interference at P (1) [accept: waves reinforce] So waves are in phase (1) [Accept: phase difference = $2\pi n$ etc] II. $S_1P - S_2P = n\lambda$ [for $n = 0, \pm 1, \pm 2, \dots$] (1) [$n = 0$ for central maximum, $n = 1$ for next one out from centre], $n = 2$ at P . (1) So $S_1P - S_2P = 0.024$ m (1) [Geometric method based upon Pythagoras ✓✓✓ if correct]	2 3
		(b) Interpose a grille of parallel metal rods and rotate. (1) The signal strength varies. (1) [Accept rotation of the sensor / aerial]	2
(c) Any 2 × (1) of: <ul style="list-style-type: none"> • the radiation penetrates the potato ✓ • absorbed within the potato, heating interior ✓ • waves transfer energy [or equiv] ✓ • water content heated / water molecules made to vibrate more ✓ 	2		
			[15]

Question		Marking details	Marks Available	
3	(a)	 <p>Correctly drawn ray (1) c shown correctly (1)</p>	2	
	(b)	$1.520 \sin \theta_A = 1.550 \sin \theta_B$ (1) [or by impl.] $\theta_A = 90^\circ$, $\theta_B = c$ (1) [or by impl.] $c = 79^\circ$ (1)	3	
	(c)	(i) 11° [$\pm 1^\circ$] e.c.f. (1)	1	
	(d)	(ii) Some enters the cladding (1) and is lost (1) Some is reflected but lost on subsequent reflections (1). (3)	3	
		Paths at different angles to the axis are of different lengths (1). Data travelling on different paths arrive different times [or by clear implic.](1) so data is muddled / smeared out / data pulses overlap (1) (3)	3	
			[12]	
4.	(a)	(i)	Photons hit the caesium surface. (1) Electrons knocked out (1) <ul style="list-style-type: none"> • Electrons cross vacuum to collecting electrode ✓ • returned to the caesium via cell and meter ✓ • constituting an electric current ✓ • aided by [p.d. of] cell ✓ } any 1 ×	3
		(ii)	Larger current (1) because more photons arrive [per second] (1)	2
	(b)	(i)	<ul style="list-style-type: none"> • Power supply polarity needs reversing ✓ • Voltage needs to be variable ✓ • voltmeter needed ✓ } any 2 ×	2
		(ii)	$E_{k \max} = 6.6 \times 10^{-34} \times 8.6 \times 10^{14} - 3.1 \times 10^{-19} \text{ J}$ (1) $= 2.6 \times 10^{-19} \text{ J}$ (1)	2
	(iii)	$E_k = \frac{1}{2}mv^2$ <u>with</u> $m = 9.1 \times 10^{-31} \text{ kg}$ (1) Convincing substitution of $v = 7.5 \times 10^5 \text{ m s}^{-1}$ to obtain $E_k = 2.6 \times 10^{-19} \text{ J}$ or vice versa (1)	2	
	(iv)	Intensity doesn't affect individual photon energies [or equiv.] (1)	1	
				[12]

Question		Marking details	Marks Available	
5	(a)	(i) $\Delta E = \frac{hc}{\lambda}$ [or $\Delta E = hf$ and $f = \frac{c}{\lambda}$] [or by impl.] (1) $\Delta E = 1.9 \times 10^{-19}$ J [or by impl.] (1) $\lambda = 1.0 \times 10^{-6}$ m (1) ((unit))	3	
		(ii) infrared	1	
		(iii) [Incident] photon causes emission of a photon (1) + 2 × (1) of: <ul style="list-style-type: none"> Incident photon energy needs to be $E_A - E_B$ [or equiv.] ✓ Emitted photon has same energy (or λ or f) as incident photon. ✓ Emitted photon in phase with incident photon. ✓ 	3	
		(iv) Two photons where there was one before [and the process repeats]	1	
	(b)	(i) More electrons in level A than in level B.	1	
		(ii) If more electrons in B than A, absorption of photons is more likely than stimulated emission.	1	
		(iii) B almost empty [because electrons ‘fall’ from B to ground state] (1) So not many electrons needed in A to cause population inversion. (1)	2	
				[12]
	6.	(a)	Weak (1) because neutrinos only feel the weak force [as well as gravity] (1) [Or because the weak force alone can cause a change of quark type].	2
		(b)	(i) Ar has 1 more proton than Cl, but electron also appears [so net charge is conserved]. [Or Ar appears as + ion (and picks up an electron)]	1
(ii) ν_e on left is a lepton [or has a lepton number of 1]; electron on right is a lepton [or]			1	
(c)		(i) 20 (ii) 19 [both answers correct]	1	
(d)		(i) udd	1	
		(ii) In version at top, neutron is lost and proton is gained. (1) [or $n + \nu_e \rightarrow p + e^-$] We can regard this as a neutron losing a d [quark] and gaining a u [quark] (1)	2	
			[8]	

Question		Marking details	Marks Available
7	(a)	[A body with a surface that] absorbs all radiation[accept: 'light'] falling upon it.	1
	(b)	(i) Area of sphere of radius $8.1 \times 10^{16} \text{ m} = 4\pi \times (8.1 \times 10^{16})^2$ (1) $[= 8.2 \times 10^{34} \text{ m}^2]$ Power reaching surface = $1.2 \times 10^{-7} \times 4\pi \times (8.1 \times 10^{16})^2$ W (1) [Or reverse argument from power to intensity, if clear] e.c.f on numerical factors in area [not for use of 2π]	2
		(ii) Absorption / scattering [of radiation by interstellar dust / gas]	1
		(iii) $9.9 \times 10^{27} = 5.67 \times 10^{-8} A \times 9900^4$ [or by impl.] (1) (Data subst. at any stage) Transposition at any stage (1) $r = 1.2 \times 10^9 \text{ m}$ (1) [e.c.f. on A , if π^2 used]	3
		(iv) Curve of correct general shape sketched which is <ul style="list-style-type: none"> • lower throughout (1) • has a maximum at longer λ (1) 	2
	(c)	Atoms / ion / [accept molecules] of a star's atmosphere (1) [or interstellar space or Earth's atmosphere] absorb specific wavelengths (1) [from the continuous spectrum] promoting electrons to higher energy level (1) [or re-emitting in all directions]	3
			[12]